



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

cases of importation. He calls attention to the fact that *Sphaerotheca tomentosa* (Otth.) Jacz.—of which *Erysiphe gigantasca* Sorok., reported on *Euphorbia virgata* from Moscow, is a synonym—is somewhat different from *Sphaerotheca mors-uvæ* (Schw.) with which Salmon in his Monograph of the Erysiphaceae, unites it. Magnus is therefore of the opinion that the form on *Euphorbia* is really a different species, saying moreover “bei der verschiedenen geographischen Verbeitung beider Formen zweifle ich nicht, dass sie zwei verschiedenen Arten entsprechen.”

PROFESSOR P. HENNINGS has published in *Hedwigia*, March, 1902, under the title, “Fungi blumenavienses II, a cl. Alfr. Möller lecti,” an article of 33 pages, containing an enumeration of the Ascomycetes from Blumenau (South Brazil) obtained in 1891-3. Some of the species had been previously reported, but are also included in this article. They are all grouped in proper systematic order under Pyrenomycetes and Discomycetes, 8 families of the former and 12 of the latter. The new genera proposed are *Aschersoniopsis* (Hypocreaceae), *Moelleroclavus* (Xylariaceae), *Stilbohypoxyton* (Xylariaceae), *Midotiopsis* (Dermateaceae), *Bulgariopsis* (Bulgariaceae), and *Moellerodiscus* (Cudoniaceae). Seventy-two new species are described, the diagnoses in Latin.

OHIO FUNGI EXSICCATI.

The fifth fascicle of the Ohio Fungi is today issued. It completes a century—the fascicles averaging 20 specimens. The 100 specimens belong to the following genera: *Aecidium* (8 sp.), *Albugo* (1 sp.), *Cercospora* (2 sp.), *Cintractia* (2 sp.), *Exoascus* (1 sp.), *Gymnoconia* (1 sp.), *Gymnosporangium* (2 sp.), *Melampsora* (2 sp.), *Microsphaera* (1 sp.), *Peronospora* (1 sp.), *Piggotia* (1 sp.), *Phyllachora* (2 sp.), *Phyllosticta* (3 sp.), *Plasmopara* (1 sp.), *Polystictus* (1 sp.), *Puccinia* (25 sp.), *Rhytisma* (1 sp.), *Septoria* (6 sp.), *Urocystis* (2 sp.), *Uromyces* (8 sp.), *Ustilago* (4 sp.), *Venturia* (1 sp.). Although instituted for the purpose of exchange with botanists many requests for the purchase of the specimens suggested at once the enlargement of the edition; now all can be accommodated who may wish the fascicles as they are issued. The price fixed, intended to cover but part of the actual cost, is \$5.00 for each five fascicles (100 specimens). Four or five fascicles will probably appear during the coming year, the labels as heretofore having a reprint of the original description of the species.